



Malvin Z-chalcone: An unexpected new open cavity for the ferric cation

Florian George, Paulo Figueiredo, Raymond Brouillard*

Laboratoire de Chimie des Polyphénols, UMR 7509 du C.N.R.S., Faculté de Chimie, Université Louis Pasteur, 1, rue Blaise Pascal, 67008 Strasbourg, France

Received 1 April 1998

Abstract

It is demonstrated that complexation between the ferric cation and the Z-chalcone of the naturally occurring anthocyanin malvin takes place in acidic aqueous solutions. The flexible open cavity of the Z-chalcone best fits the steric and electronic requirements of the ferric ion in water. © 1999 Published by Elsevier Science Ltd. All rights reserved.

Keywords: Malvin chloride; Ferric chloride; Z-chalcone complexation; Molecular modelisation

1. Introduction

Among the many metallic species occurring in plant biological systems and known to interact with anthocyanins, only aluminium (III) was thoroughly studied (Harborne, 1989; Elhabiri, Figueiredo, Toki, Saito & Brouillard, 1997). Moreover, in the anthocyanin family, only those members possessing an *ortho*-dihydroxy aromatic moiety or catechol group in their B ring, like cyanin for instance, covalently bind to trivalent cations such as Al^{3+} and Ga^{3+} , as it has been regularly demonstrated over the past three decades with aluminium (Elhabiri et al., 1997; Asen, Norris & Stewart, 1969; Takeda, 1977). Aluminium–anthocyanin complex formation provides colour stabilization and variation to the pigment containing media.


Iron being one of the most important factors for the growth of plant species, this prompted us to investigate the possible interaction between iron (III) and the flavylium form of some anthocyanins with a catechol moiety. The result was surprising and, at first sight, rather disappointing: instead of the strong colour stabilization and variation seen with Al^{3+} , loss of col-

our was observed. We repeated the experiment using an anthocyanin without a catechol group, malvin chloride (Scheme 1) and again the presence of the ferric cation resulted in loss of colour. We came thus to the assumption that the Fe^{3+} /anthocyanin bleaching effect had little to do with the presence or the absence of a catechol group in the anthocyanin. It is the aim of the present work to establish the mechanism accounting for the bleaching of malvin aqueous solutions in the presence of ferric chloride.

2. Results and discussion

Malvin and ferric chlorides were made to react at 25°C and an equilibrium state was attained after approximately 17 h (Fig. 1). From Fig. 1, two spectroscopic features emerge. One is the presence of two isosbestic points at 284 and 362 nm, which strongly suggests the existence of a chemical equilibrium, apparently between the flavylium cation, the only malvin stable form at pH 1, and an unknown molecular species involving Fe^{3+} . The other is a steady decrease in the intensity of the flavylium visible absorption band (colour loss) whose maximum wavelength (518 nm) remains remarkably constant. This last obser-

* Author to whom correspondence should be addressed.



Scheme 1.

vation allows one to eliminate the flavylum cation as being the malvin form interacting with ferric chloride.

Subtraction of flavylum and ferric ions absorption contributions from the global UV-visible absorption spectra (Fig. 1) leads to the spectra shown in Fig. 2. The result indicates that at pH 1 a new species, with a maximum of absorption around 340 nm, appears. Moreover, when chlorogenic acid, a good malvin copigment (Mazza & Brouillard, 1990), was initially added to a solution containing malvin and ferric chlorides, the reaction did not take place. Thus, it can be concluded that covalent hydration of the flavylum form is a key step in the appearance of the malvin form interacting with Fe^{3+} in water. Kinetic and spectral characteristics of malvin structural transformations (Markham, 1982; Brouillard & Lang, 1990; Santos, Turner, Lima, Figueiredo, Pina & Maçanita, 1993) suggest that the species interacting with the ferric ion is the Z-chalcone (Scheme 1).

Confirmation of that unexpected and somewhat extraordinary result was given by monitoring the reaction with the help of ^1H NMR spectroscopy. Fig. 3(a) shows a spectrum of a pure malvin solution, at 25°C

in $\text{DCl}/\text{D}_2\text{O}$ with a pD value close to 1.50. In it, can be observed the peaks corresponding to the five malvin forms at equilibrium in slightly acidic to acidic aqueous solutions (Brouillard & Lang, 1993; Santos et al., 1993). Adding a large excess of ferric chloride, at constant pD, to the initial malvin solution leads to the simplification over time, of the “complex” initial spectrum. For instance, after 1 h [Fig. 3(b)], the two hemiacetals and the E-chalcone (Brouillard & Lang, 1993; Santos et al., 1993) are still present but a new species now exists in the solution while the flavylum ion and the free Z-chalcone have practically disappeared from the medium. Finally, after complete evolution of the system toward equilibrium, NMR spectroscopy indicates that the only stable species remaining in the solution is the Z-chalcone complex formed when FeCl_3 is added to the solution [Fig. 3(c)].

The ^1H NMR experiment is in good agreement with and completes what was observed using UV-visible absorption spectroscopy: Fe^{3+} associates selectively with only one of the five malvin free forms and this is the Z-chalcone. In the aromatic range significant features are: H-2' and H-6' are no longer equivalent; H-




Fig. 1. UV-visible absorption spectra, as a function of time, of malvin chloride at pH 1, in the presence of an excess of ferric chloride. $T = 25^\circ\text{C}$.




Fig. 2. Spectra deduced from the spectra shown on Fig. 1 by subtraction of the flavylum and the ferric ions absorption contributions. $T = 25^\circ\text{C}$.




Fig. 3. ^1H NMR spectra of malvin chloride without FeCl_3 (a). ^1H NMR spectra of malvin chloride with FeCl_3 : after 1 h (b) and at equilibrium after 1020 min (c). The peaks of the different malvin forms are indicated with the following symbols: (*) flavylum cation; (○/●) R/S hemiacetal; (▲) E-chalcone; (◆) Z-chalcone and (□) Z-chalcone-iron complex. $T = 25^\circ\text{C}$.

$2'$ is shielded and H-6' is deshielded. H-6 and H-8 are still observed while H-4 is no longer observable. In the absence of Fe^{3+} , the transversal T_2 relaxation time values range from 1 to 2 s. Introduction of the ferric ion lowers all those values down to *ca.* 0.3 s. This may account for an iron anisotropic effect, which might explain why H-4 is removed from the spectrum.

The 3D structure of the new iron complex (Fig. 4) was visualized through molecular modelling using the Hyperchem software in the ZINDO/1 semi-empirical mode (Zerner, 1991). Parameters were taken from comparisons between the properties of various flavylum ions studied in our laboratory and a few crystallographic data available from the literature (Ueno & Saito, 1977a,b). ZINDO/1 results demonstrate that the ferric ion is positioned between the B ring of the chro-

mophore and the glucosyl residue at C-5 of the aglycone (Fig. 4). Under the Fe^{3+} effect, it is thought that the H-4 signal has been enlarged over an extent that makes it impossible to be observed within the window of our experiment. Moreover, the splitting of the signal attributed to protons H-2' and H-6' into two well separated singlets may also be explained by a conformation where the B ring rotation is precluded. The presence of the ferric ion within the Z-chalcone open cavity induces another interesting feature highly stabilizing the Z-chalcone complex. This is the existence of two hydrogen bonds, one between the hydroxyl at C-9 and the oxygen at C-3 and the other between the oxygen at C-9 (acceptor) and the hydroxyl at C-2 of the 3-glucosyl residue (Fig. 4). No such hydrogen bonds are to be found in the ZINDO/1 model of the free Z-chalcone (Fig. 4). We also noted that for 3-deoxyanthocyanidins and from UV-visible observations, iron (III) complexation does not take place. This result points to the importance of the unusual H-bonds found in the model of the malvin complex.

More work is under progress in our laboratory where we are investigating many systems made of natural anthocyanins devoid of a catechol moiety and exposed to ferric chloride. Preliminary results are extremely encouraging. Iron is a common metal in living systems frequently carried by siderophores (Blanc et al., 1997). In the case of the malvin Z-chalcone ferric complex, we found for its complexation constant a value of $\approx 10^5$ (aqueous medium). This is larger than the values determined for siderophores bearing catechol groups like the enterobactin iron (III) complex (Avdeef, Sofen, Bregante & Raymond, 1978; Harris et al., 1979). The important role that anthocyanins




Fig. 4. Semi-empirical modelisation in the ZINDO/1 parametrization of the Z-chalcone iron complex (left) and the free Z-chalcone.

could play as antioxidants in living systems may well be explained by their possible regulation of iron uptake and activity within the most complicated biological media. Could the *raison d'être* of anthocyanins inside fruits, tubers and roots for instance, be related to this until now unknown biological function?

3. Experimental

3.1. Materials

Malvin chloride (99% pure) was purchased from Roth and ferric chloride hexahydrate from Aldrich. Both were used as supplied.

3.2. Absorption spectra

Absorption spectra were recorded with a Hewlett Packard diode-array spectrophotometer fitted with a quartz cell ($d = 1$ cm) equipped with a stirring magnet. A constant temperature of $25 \pm 0.1^\circ\text{C}$, measured with a Comark thermocouple, was maintained in the spectrometer sample cell by use of a Lauda water-thermostated external bath. Water used in sample preparations was bidistilled.

3.3. Data analysis

UV-visible absorption data were recorded on a Pentium P200 PC using the Hewlett Packard UV-visible Chemstation Programme. Mathematical treatments applied to the data were carried out with the Microsoft Excel 5.0 programme running on a PowerMacintosh PM7200/90 machine.

3.4. Kinetic measurements

The mother solution of malvin chloride (1×10^{-4} M) was prepared in 0.10 M hydrochloric acid and left to equilibrate in the dark for about 2 h. 1 ml of this solution was put into the spectrometer sample cell. Then to it was quickly added 1 ml of ferric chloride (1×10^{-3} M), also prepared with 0.10 M hydrochloric acid. Complete UV-visible spectra were immediately recorded every 10 min until final equilibrium was reached. Some are shown in Fig. 1.

3.5. ^1H NMR spectra

Malvin chloride was dissolved in DCl/D₂O (*ca.* 0.1 M) to a concentration of 1.5×10^{-4} M and a pD value of 1.5. The ferric chloride initial solution

(1×10^{-2} M) was also brought to the same free acidity value using DCl in D₂O. ^1H NMR spectra were run on a solution obtained by mixing given volumes of the two above solutions. Spectra were recorded with a Bruker AMX-500 spectrometer.

Assignments of the observed chemical shifts to the different aromatic protons of each species were made according to (Santos *et al.*, 1993). δ/ppm : [(*) flavylum cation] 8.97 (1H, *s*, H-4), 7.82 (2H, *s*, H-2' and H-6'), 7.10 (1H, *s*, H-8), 7.00 (1H, *s*, H-6); [(○/●) *R/S* hemiacetal] 6.98/6.95 (2H, *s*, H-2' and H-6'), 6.60/6.58 (1H, *s*, H-4), 6.44 (1H, *s*, H-6), 6.25 (1H, *s*, H-8); [(▲) *E*-chalcone] 7.20 (2H, *s*, H-2' and H-6'), 6.70 (1H, *s*, H-4), 6.04 (1H, *s*, H-6), 5.87 (1H, *s*, H-8); [(◆) *Z*-chalcone] 7.33 (2H, *s*, H-2' and H-6'), 6.80 (1H, *s*, H-4), 6.32 (1H, *s*, H-6), 6.22 (1H, *s*, H-8); [(□) complex] 7.79 (1H, *s*, H-6'), 6.67 (1H, *s*, H-8), 6.61 (1H, *s*, H-6), 5.95 (1H, *s*, H-2').

3.6. Semi-empirical quantum mechanical calculations

Semi-empirical quantum mechanical calculations were performed on a Pentium P200 PC using the Hyperchem programme (version 4, Hypercube, Ont., Canada) in the MM+ and ZINDO/1 parametrizations.

References

- Asen, S., Norris, K. H., & Stewart, R. N. (1969). *Phytochemistry*, 8, 653.
- Avdeef, A. E., Sofen, S. R., Bregante, T. L., & Raymond, K. N. (1978). *J. Am. Chem. Soc.*, 100, 5362.
- Blanc, S., Yakirevitch, P., Leize, E., Meyer, M., Libman, J., Vandorsselaer, A., Albrecht-Gary, A. M., & Shanzer, A. (1997). *J. Am. Chem. Soc.*, 119, 4934.
- Brouillard, R., & Lang, J. (1990). *Can. J. Chem.*, 68, 755.
- Elhabiri, M., Figueiredo, P., Toki, K., Saito, N. & Brouillard, R. (1997). *J. Chem. Soc. Perkin Trans. II*, 355.
- Harborne, J. B. (1989). In J. B. Harborne, *Methods in plant biochemistry*, Vol. 1 (pp. 1–28). London: Academic Press.
- Harris, W. R., Carrano, C. J., Cooper, S. R., Sofen, S. R., Avdeef, A. E., McArdle, J. V., & Raymond, K. N. (1979). *J. Am. Chem. Soc.*, 101, 6097.
- Markham, K. R., *Techniques of flavonoid identification*. London, Academic Press, 1982.
- Mazza, G., & Brouillard, R. (1990). *Phytochemistry*, 29, 1097.
- Santos, H., Turner, D. L., Lima, J. C., Figueiredo, P., Pina, F. S., & Maçanita, A. L. (1993). *Phytochemistry*, 33, 1227.
- Takeda, K. (1977). *Proc. Jpn. Acad. B*, 53, 257.
- Ueno, K., & Saito, N. (1977). *Acta Cryst. B*, 33, 111.
- Ueno, K., & Saito, N. (1977). *Acta Cryst. B*, 33, 114.
- Zerner, M. C. (1991). In K. B. Lipkowitz and D. B. Boyd. *Reviews in Computational chemistry*, vol. 2 (pp. 313–366). New York: VCH Publishers.